



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> : C07K 15/04, C12N 15/62	A1	(11) International Publication Number: WO 95/00555 (43) International Publication Date: 5 January 1995 (05.01.95)
<p>(21) International Application Number: PCT/EP94/02088</p> <p>(22) International Filing Date: 28 June 1994 (28.06.94)</p> <p>(30) Priority Data: 93110298.2 28 June 1993 (28.06.93) EP</p> <p>(34) Countries for which the regional or international application was filed: DE et al.</p> <p>(71) Applicant (for all designated States except US): EUROPEAN MOLECULAR BIOLOGY LABORATORY (EMBL) [DE/DE]; Meyerhofstrasse 1, D-69117 Heidelberg (DE).</p> <p>(72) Inventor; and</p> <p>(75) Inventor/Applicant (for US only): STEWART, Francis [AU/DE]; Pfarrgasse 7, D-69251 Gaiberg (DE).</p> <p>(74) Agents: WEICKMANN, H. et al.; Kopernikusstrasse 9, D-81679 München (DE).</p>	<p>(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b></p> <p><i>With international search report.</i></p> <p><i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p> <p><i>With an indication in relation to a deposited microorganism furnished under Rule 13bis separately from the description.</i></p> <p><i>Date of receipt by the International Bureau:</i> 23 September 1994 (23.09.94)</p>	
<p>(54) Title: REGULATION OF SITE-SPECIFIC RECOMBINATION BY SITE-SPECIFIC RECOMBINASE/NUCLEAR RECEPTOR FUSION PROTEINS</p>		
<p>(57) Abstract</p> <p>An invention is described that permits the regulation of recombination in cells, organisms or appropriate cell-free systems. The invention involves creating fusion proteins between recombinase proteins, or components of recombinase systems, and ligand binding domains derived from nuclear receptors. The fusion proteins show little recombinase activity in the absence of the ligand that binds to the ligand binding domain. Upon binding of the ligand, recombinase activity is induced. The invention provides a practical means to regulate recombination in cells and organisms and, by linking ligand binding to recombination, provides a simple means whereby ligand binding can be measured as recombination achieved.</p>		

FOR THE PURPOSES OF INFORMATION ONLY					
Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.					
AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CH	Switzerland	KR	Republic of Korea	SG	Singapore
CN	China	KZ	Kazakhstan	SI	Slovenia
CO	Colombia	LA	Laos	SK	Slovakia
CR	Costa Rica	LB	Lebanon	SM	San Marino
CU	Cuba	LU	Luxembourg	SN	Senegal
CZ	Czech Republic	LV	Latvia	SR	Sri Lanka
DE	Germany	MC	Monaco	SS	South Sudan
DK	Denmark	MD	Republic of Moldova	ST	Sao Tome and Principe
ES	Spain	MG	Madagascar	TD	Chad
FI	Finland	MH	Maldives	TE	Togo
FR	France	MI	Mali	TJ	Tajikistan
GA	Gabon	ML	Moldova	TT	Trinidad and Tobago
GB	United Kingdom	MM	Myanmar	UA	Ukraine
GD	Grenada	MO	Macao	US	United States of America
GE	Georgia	MP	Marshall Islands	UZ	Uzbekistan
GG	Guernsey	MR	Mauritania	VC	St. Vincent and the Grenadines
GH	Ghana	MT	Malta	VG	Virgin Islands, British
GI	Guinea-Bissau	MU	Mauritius	VI	Virgin Islands, U.S.
GL	Greenland	NB	Norway	WS	Samoa
GN	Guinea	NC	New Caledonia		
GR	Greece	NE	Niger		
GT	Guatemala	NL	Netherlands		
GU	Guam	NO	Norway		
GW	Guinea-Bissau	NZ	New Zealand		
GY	Guyana	PL	Poland		
HA	Haiti	PT	Portugal		
HN	Honduras	RO	Romania		
HR	Croatia	RU	Russian Federation		
HT	Haiti	SD	Sudan		
HU	Hungary	SG	Singapore		
IE	Ireland	SI	Slovenia		
IL	Israel	SK	Slovakia		
IN	India	SM	San Marino		
IQ	Iraq	SN	Senegal		
IR	Iran, Islamic Republic of	SR	Sri Lanka		
IS	Iceland	SS	South Sudan		
IT	Italy	ST	Sao Tome and Principe		
JE	Jersey	TD	Chad		
JM	Jamaica	TE	Togo		
JO	Jordan	TJ	Tajikistan		
JP	Japan	TK	Tokelau		
KE	Kenya	TL	Timor-Leste		
KG	Kyrgyzstan	TM	Turkmenistan		
KH	Kampuchea	TN	Tunisia		
KI	Kiribati	TR	Turkey		
KM	Comoros	TV	Tuvalu		
KN	Saint Kitts and Nevis	UG	Uganda		
KR	Republic of Korea	US	United States of America		
KU	Kuwait	UZ	Uzbekistan		
KV	Kiribati	VC	St. Vincent and the Grenadines		
KW	Kuwait	VE	Venezuela		
KY	Cayman Islands	VG	Virgin Islands, British		
KZ	Kazakhstan	VI	Virgin Islands, U.S.		
LA	Laos	WS	Samoa		
LB	Lebanon				
LC	St. Lucia				
LI	Liechtenstein				
LK	Sri Lanka				
LU	Luxembourg				
LV	Latvia				
LY	Libya				
MC	Monaco				
MD	Republic of Moldova				
ME	Montenegro				
MG	Madagascar				
MH	Marshall Islands				
MI	Mali				
ML	Mali				
MM	Myanmar				
MO	Macao				
MP	Marshall Islands				
MR	Mauritania				

FOR THE PURPOSES OF INFORMATION ONLY					
Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.					
AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CH	Switzerland	KR	Republic of Korea	SG	Singapore
CN	China	KZ	Kazakhstan	SI	Slovenia
CO	Colombia	LA	Laos	SK	Slovakia
CR	Costa Rica	LB	Lebanon	SM	San Marino
CU	Cuba	LU	Luxembourg	SN	Senegal
CZ	Czech Republic	LV	Latvia	SR	Sri Lanka
DE	Germany	MC	Monaco	SS	South Sudan
DK	Denmark	MD	Republic of Moldova	ST	Sao Tome and Principe
ES	Spain	MG	Madagascar	TD	Chad
FI	Finland	MH	Maldives	TE	Togo
FR	France	ML	Mali	TJ	Tajikistan
GA	Gabon	MN	Mongolia	TT	Trinidad and Tobago
GB	United Kingdom	MO	Macao	UA	Ukraine
GD	Grenada	MP	Marshall Islands	US	United States of America
GE	Georgia	MR	Mauritania	UZ	Uzbekistan
GG	Guernsey	MT	Malta	VC	St. Vincent and the Grenadines
GH	Ghana	MU	Mauritius	VG	Virgin Islands, British
GI	Guinea-Bissau	NL	Netherlands	VI	Virgin Islands, U.S.
GL	Greenland	NO	Norway	WS	Samoa
GM	Gambia	NZ	New Zealand		
GN	Guinea	PL	Poland		
GR	Greece	PT	Portugal		
GT	Guatemala	RO	Romania		
GU	Guam	RU	Russian Federation		
GW	Guinea-Bissau	SD	Sudan		
GY	Guyana	SG	Singapore		
HA	Haiti	SI	Slovenia		
HN	Honduras	SK	Slovakia		
HR	Croatia	SM	San Marino		
HT	Haiti	SN	Senegal		
HU	Hungary	SR	Sri Lanka		
IE	Ireland	SS	South Sudan		
IL	Israel	ST	Sao Tome and Principe		
IN	India	TD	Chad		
IO	British Indian Ocean Territory	TE	Togo		
IS	Iceland	TJ	Tajikistan		
IT	Italy	TK	Turkmenistan		
JE	Jersey	TL	Timor-Leste		
JM	Jamaica	TM	Turkmenistan		
JO	Jordan	TT	Trinidad and Tobago		
JP	Japan	UA	Ukraine		
KE	Kenya	US	United States of America		
KG	Kyrgyzstan	UZ	Uzbekistan		
KH	Kampuchea	VC	St. Vincent and the Grenadines		
KI	Kiribati	VG	Virgin Islands, British		
KM	Comoros	VI	Virgin Islands, U.S.		
KN	St. Kitts and Nevis	WS	Samoa		
KO	North Korea				
KR	South Korea				
KU	Kuwait				
KV	Kingdom of the Netherlands				
KZ	Kazakhstan				
LA	Laos				
LB	Lebanon				
LC	St. Lucia				
LI	Liechtenstein				
LK	Sri Lanka				
LU	Luxembourg				
LV	Latvia				
LY	Libya				
MC	Monaco				
MD	Republic of Moldova				
ME	Montenegro				
MG	Madagascar				
MH	Marshall Islands				
MI	Mali				
MJ	Maldives				
MK	Macedonia				
ML	Mali				
MM	Myanmar				
MN	Mongolia				
MO	Macao				
MP	Marshall Islands				

FOR THE PURPOSES OF INFORMATION ONLY					
Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.					
AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CH	Switzerland	KR	Republic of Korea	SG	Singapore
CN	China	KZ	Kazakhstan	SI	Slovenia
CO	Colombia	LA	Laos	SK	Slovakia
CR	Costa Rica	LB	Lebanon	SM	San Marino
CU	Cuba	LU	Luxembourg	SN	Senegal
CZ	Czech Republic	LV	Latvia	SR	Sri Lanka
DE	Germany	MC	Monaco	SS	South Sudan
DK	Denmark	MD	Republic of Moldova	ST	Sao Tome and Principe
ES	Spain	MG	Madagascar	TD	Chad
FI	Finland	MH	Maldives	TE	Togo
FR	France	ML	Mali	TJ	Tajikistan
GA	Gabon	MN	Mongolia	TT	Trinidad and Tobago
GB	United Kingdom	MO	Macao	UA	Ukraine
GD	Grenada	MP	Marshall Islands	US	United States of America
GE	Georgia	MR	Mauritania	UZ	Uzbekistan
GG	Guernsey	MT	Malta	VC	St. Vincent and the Grenadines
GH	Ghana	MU	Mauritius	VG	Virgin Islands, British
GI	Guinea-Bissau	NL	Netherlands	VI	Virgin Islands, U.S.
GL	Greenland	NO	Norway	WS	Samoa
GM	Gambia	NZ	New Zealand		
GN	Guinea	PL	Poland		
GR	Greece	PT	Portugal		
GT	Guatemala	RO	Romania		
GU	Guam	RU	Russian Federation		
GW	Guinea-Bissau	SD	Sudan		
GY	Guyana	SG	Singapore		
HA	Haiti	SI	Slovenia		
HN	Honduras	SK	Slovakia		
HR	Croatia	SM	San Marino		
HT	Haiti	SN	Senegal		
HU	Hungary	SR	Sri Lanka		
IE	Ireland	SS	South Sudan		
IL	Israel	ST	Sao Tome and Principe		
IN	India	TD	Chad		
IO	British Indian Ocean Territory	TE	Togo		
IS	Iceland	TJ	Tajikistan		
IT	Italy	TK	Turkmenistan		
JE	Jersey	TL	Timor-Leste		
JM	Jamaica	TM	Turkmenistan		
JO	Jordan	TT	Trinidad and Tobago		
JP	Japan	UA	Ukraine		
KE	Kenya	US	United States of America		
KG	Kyrgyzstan	UZ	Uzbekistan		
KH	Kampuchea	VC	St. Vincent and the Grenadines		
KI	Kiribati	VG	Virgin Islands, British		
KM	Comoros	VI	Virgin Islands, U.S.		
KN	St. Kitts and Nevis	WS	Samoa		
KO	North Korea				
KR	South Korea				
KU	Kuwait				
KV	Kingdom of the Netherlands				
KZ	Kazakhstan				
LA	Laos				
LB	Lebanon				
LC	St. Lucia				
LI	Liechtenstein				
LK	Sri Lanka				
LU	Luxembourg				
LV	Latvia				
LY	Libya				
MC	Monaco				
MD	Republic of Moldova				
ME	Montenegro				
MG	Madagascar				
MH	Marshall Islands				
MI	Mali				
MJ	Maldives				
MK	Macedonia				
ML	Mali				
MM	Myanmar				
MN	Mongolia				
MO	Macao				
MP	Marshall Islands				

Regulation of site-specific recombination by site-specific  
recombinase/nuclear receptor fusion proteins  
**Specification**

The present invention relates to the technical field of genetic manipulation, and more specifically, to the use of recombination-mediated DNA rearrangements and to the use of the regulatory potential of nuclear receptors.

The use of site-specific recombinases (SSRs) to induce defined rearrangements of DNA has been described in a variety of organisms (1-12). These reports describe the introduction of a DNA construct that contains SSR target sites. Subsequent exposure to the SSR enzyme activity resulted in the DNA rearrangement determined by the disposition of the target sites (see reference 13 for a recent review of SSRs). Three SSRs have been used in this manner to date; FLP recombinase from the 2 $\mu$  episome of *Saccharomyces cerevisiae* (1,2,5,6,9,10), CRE recombinase from the *Escherichia coli* phage P1 (3,4,8,11,12) and R recombinase from pSR1 of *Zygosaccharomyces rouxii* (7). Amongst other SSR systems relevant to the invention described here are those listed in references 13 and 14, and SSRs from *Kluyveromyces drosophilarius* (15), *Kluyveromyces waltii* (16),  $\lambda$  Int (17) and the Gin recombination system from phage Mu (18). The content of the above document is incorporated by reference.

For many applications in cells, organisms and cell-free in vitro systems, SSR induced DNA rearrangements must be regulated. Current implementation of the potential offered by SSRs is limited by the means available to regulate SSR activity. In experiments with cultured cells, unregulated SSRs have been used. For example, after introduction of SSR target sites into

ded recombination event was regulated merely by the time of introduction of an appropriate macromolecule. Amongst other

ranged target sites and the SSR and in which the recombination event can be induced after cell numbers have been expanded.

In experiments with transgenic animals, the issue of SSR regulation has been addressed by regulating the expression of an introduced SSR gene using the inducible heat-shock promoter in *Drosophila* (5) or a tissue-specific promoter in mice (11). Both approaches have limited applicability. Namely, heat-shock regulation of transgene expression is currently only useful in flies and no suitable counterpart is available for use in cell lines or vertebrates. Also, the use of a tissue-specific promoter to regulate transgene expression relies on the limited availability of suitable promoters and enhancers and the expression pattern achieved is confined to the times and places at which these tissue specific elements are active.

The problem underlying the present invention was to provide a regulated recombination system, wherein the disadvantages of the prior art are at least partially eliminated. More specifically, the problem was to provide a recombination system, wherein the recombination event can be induced independently from any tissue specific restrictions.

This patent application describes an invention that regulates SSR activity, rather than its expression. Thus any means of achieving and directing expression can be used, such as using ubiquitous or broadly active promoters and enhancers, as well as tissue specific or inducible promoters and enhancers.

One aspect of the present invention relates to a fusion protein, comprising a recombinase protein or a component of a recombinase complex, fused to part or all of a nuclear receptor, so that the amino acids that bind the ligand of said nuclear

sence of ligand binding to said ligand binding domain, and  
(b) recombinase activity is induced or altered by binding of ligand to said ligand binding domain

Preferably the recombinase activity is altered by binding of ligand to ligand binding domain by a factor of at least 10, more preferably of at least 20 and most preferably of at least 40.

The present invention resides in the regulation of SSR activity by fusing an SSR protein to the ligand binding domain (LBD) of a nuclear receptor. SSR-LBD fusion proteins comprise an amino acid sequence of SSRs physically attached to the amino acid sequence of an LBD of a nuclear receptor. That is, the invention is a fusion protein, comprising a recombinase protein, or a component of a recombinase complex, fused to the ligand binding domain of a nuclear receptor. Preferably, the recombinase protein or a component of a recombinase complex is fused to the nuclear receptor or ligand binding domain thereof by means of genetic fusion, i.e. the SSR-LBD protein is a linear genetic fusion encoded by a single nucleic acid. On the other hand, the present invention also encompasses SSR-LBD fusion proteins which are linked by different means, e.g. through a spacer molecule having reactive groups thereon, which are covalently bound to each protein domain.

Most simply, the attachment of the SSR and LBD components can be achieved by making a DNA construct that encodes for the amino acid sequence of the SSR-LBD fusion protein with the LBD encoding DNA placed in the same reading frame as the SSR encoding DNA, preferably either at the amino or carboxy termini of the SSR protein (19). More preferably, the nuclear receptor or ligand binding domain thereof is fused to the C-terminus of the recombinase protein or component of a recombinase complex. In an especially preferred embodiment of the present invention the nuclear receptor or ligand binding domain thereof is fused to

SSR-LBD fusion proteins can coexist with target sites without

ments, recombinase activity of the SSR-LBD fusion protein, in the absence of the relevant ligand, is at least 200x less active than wild type recombinase activity. Upon presenting the SSR-LBD fusion proteins with the relevant ligand, recombinase activity is induced to more than 20% of wild type, that is, equal to or greater than 40x induction. This is the first description of post-transcriptional regulation of SSRs by any means. In particular, the invention permits the propagation of both the unrearranged target sites and the SSR-LBD in living systems. More specifically:

(a) cell lines can be established that carry both the unrearranged target sites and the SSR-LBD as a prerequisite for biochemical studies or introduction into organisms. The cell lines can be homogeneous and characterised before expansion to the required quantities and subsequent induction of recombination by administration of the relevant ligand,

(b) recombination can be regulated in any experimentally manipulatable multicellular organism by administration of the relevant ligand. The SSR-LBD could be introduced to the organism either through the incorporation of cells or by direct means such as microinjection or in the genome of a viral based vector. Recombination can be induced after characterisation by administration of the relevant ligand.

The term "nuclear receptor", as used herein, refers to a molecule, preferably a protein molecule, which may be glycosylated or unglycosylated, having the abilities to bind ligand and to be incorporated into a nucleus of a cell. Specifically, the term nuclear receptor refers to those proteins that display functional or biochemical properties that are similar to the functional or biochemical properties displayed by the steroid hormone receptors with respect to ligand binding, for example,

proteins that are related by their amino acid sequence to the LBDs of the steroid hormone receptors. The paper of Laudet et al. (1988) *Science* 241: 1568-1571, is hereby incorporated by reference.

ment of nuclear receptor amino acid sequences using standard methodologies. Included in the definition of nuclear receptors used here are also those proteins not listed in Laudet et al. but which can be included using the methods employed by Laudet et al.. The term nuclear receptor also includes mutant derivatives of nuclear receptor amino acid sequences which retain sufficient relatedness to nuclear receptor amino acid sequences as to be identifiable as related using the methods employed by Laudet et al.

The nuclear receptor which is fused to the recombinase protein is preferably a hormone receptor, e.g. a receptor recognized by steroids, vitamins or related ligands. Examples of suitable nuclear receptors are listed in reference 20, which is hereby incorporated by reference. Preferably, the nuclear receptor is a steroid hormone receptor, more preferably, a glucocorticoid, estrogen, progesteron, or androgen receptor.

In the SSR-LBD fusion protein of the present invention, it is not required that the complete nuclear receptor is present; i.e. it is sufficient that the amino acids that bind the ligand are fused to the SSR.

Upon binding their relevant ligand, nuclear receptors become active, or altered, transcription factors. The cloning of cDNAs encoding members of the steroid receptor family of proteins revealed that they share amino acid sequence homology, in particular in the protein domain that binds ligand. The LBD can be separated from the rest of the protein and fused to other transcription factors conferring ligand regulation onto the resulting fusion proteins. For the glucocorticoid and estrogen receptors, the domain that binds ligand has been fused to other transcription factors and

experimentally regulated in this manner so far are also transcription factors. Transcription factors and oncoproteins are not encoded in the classical sense and the regulation of their

26, is essentially similar to the role that LBDs play in regulating steroid receptors. Prior attempts to extend the regulatory possibilities of the LBD fusion protein strategy beyond transcription factors and oncoproteins have yet to be reported. Attempts to regulate the enzyme activities of either  $\beta$ -galactosidase or dihydrofolate reductase by fusing them with the glucocorticoid LBD have been unsuccessful (D. Picard, public seminar given at EMBL, Heidelberg, Nov. 1991).

The present invention extends the regulatory possibilities of the LBD fusion protein strategy beyond the currently documented transference of ligand regulation from nuclear receptors to other transcription factors, to include an enzyme activity, namely a site specific recombinase. The term "site specific recombinase" refers to any protein component of any recombinant system that mediates DNA rearrangements in a specific DNA locus, including site specific recombinases of the integrase or resolvase/invertase classes (13; Abremski, K.E. and Hoess, R.H. (1992) Protein Engineering 5, 87-91; Khan, E., Mack, J.P.G., Katz, R.A., Kulkosky, J. and Skalka, A.M. (1991) Nucleic acids Res. 19, 851-860) and site-specific recombination mediated by intron-encoded endonucleases (Perrin, A., Buckle, M. and Dujon, B. (1993) EMBO J. 12, 2939-2947). Preferred recombinase proteins, which can be used in the fusion proteins according to the invention, are selected from the group consisting of: FLP recombinase, Cre recombinase, R recombinase from the *Zygosaccharomyces rouxii* plasmid pSR1, A recombinase from the *Kluyveromyces drosophilarius* plasmid pKD1, A recombinase from the *Kluyveromyces waltii* plasmid pKW1, any component of the  $\lambda$  Int recombination system, any component of the Gin recombination system, or variants thereof. The term "variant" in this context refers to proteins which are derived from the above proteins by

could retain the ability to act as a recombinase, or it could retain protein/protein or protein/DNA interactions critical to



In a preferred embodiment of the invention, FLP recombinase is fused to the LBD of the estrogen, glucocorticoid, progesterone or androgen receptors (20). Other preferred embodiments include fusing Cre recombinase or R recombinase, or SSRs from *Kluyveromyces drosophilarius* or *Kluyveromyces waltii*, to these LBDs. Another preferred embodiment involves regulating one or more components of an SSR complex to these LBDs, in particular, components of the  $\lambda$  Int or Gin recombination systems. The invention, in providing a means to regulate recombination, is however not limited to known recombinases and recombination complexes and is not limited to known nuclear receptor LBDs. Rather, the strategy of fusing recombinases, or components of recombination complexes, to LBDs of nuclear receptors is applicable to any fusion combination of these proteins which display the desired characteristics readily identifiable without undue experimentation on the part of a skilled person.

A further subject-matter of the present invention is a nucleic acid which encodes the fusion protein according to the present invention. Preferably, the nucleic acid is a DNA or RNA.

Still a further subject-matter of the present invention is a recombinant vector comprising at least one copy of the nucleic acid as defined above. This recombinant vector may be a eukaryotic vector, a viral vector, or a prokaryotic vector, or a vector which can be maintained in eukaryotic and prokaryotic host cells. The recombinant vector is obtainable by inserting a nucleic acid encoding a SSR-LBD fusion protein into a suitable starting vector. Specific examples of suitable starting vectors are given for example in Molecular Cloning. A Laboratory Manual, 2nd edition, J. Sambrook et al. (1989), Cold Spring Harbor Laboratory Press, chapters 1, 2, 3, 4, 16 and 17.

Extra-chromosomal vectors, e.g. circular plasmids are especially preferred. An example of a plasmid vector is the plasmid pHEE1 comprising a nucleic acid encoding a fusion of

gen receptor. The plasmid pHFE1 has been deposited at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ), Mascheroder Weg 1b, D-38124 Braunschweig on June 25, 1994 under the accession number DSM 9265 according to the requirements of the Budapest Treaty.

Still a further subject-matter of the present invention is a cell containing a nucleic acid or a recombinant vector as defined above. Preferably, the cell is stably transformed with the nucleic acid or the vector. Suitable cells are eukaryotic or prokaryotic cells. Examples of prokaryotic cells are gram-negative bacterial cells, especially E.coli cells. Examples of eukaryotic cells are mammalian cells, yeasts and trypanosomes.

The invention separates the function of ligand binding from the functions of transcription factors, coupling ligand binding to recombinase activity. Thereby, the binding of ligand can be assessed by any means that measures recombinase activity. Thus, further embodiments of the invention include methods for determining the binding of ligand to the LBD of a nuclear receptor, comprising the steps of:

(i) the introduction of the SSR-LBD fusion protein or the nucleic acid coding therefor into cells, or appropriate cell-free systems, that contain the DNA target sites for the SSR; (ii) administering the ligand or a mixture suspected to contain a ligand or ligands to be evaluated; if the ligand is not already present in the system; (iii) detecting the recombinase activity, if any, of the SSR-LBD by detecting, directly or indirectly, recombination or changes in the recombination rate between the DNA target sites.

The introduction of SSR-LBD fusion proteins into cells may be accomplished by transforming cells with a vector containing the SSR-LBD fusion protein.

The invention is particularly useful in systems which already contain the ligand and in which the ligand-concentration is determined by detecting recombination between DNA target sites.

Particularly preferred embodiments include:

- (a) direct measurements of the physical arrangement of the target sites using techniques such as gel electrophoresis of DNA molecules, Southern blotting or polymerase chain reaction.
- (b) indirect measurements by assessing the properties encoded by the DNA regions carrying the target sites before or after recombination. For example, recombination could excise an antibiotic resistance gene from the genome of the host and thus recombination can be measured as sensitivity to the antibiotic.
- (c) the measurement of intracellular ligand concentrations by evaluating recombination between the DNA target sites resident in the cells that also carry the SSR-LBD fusion proteins.
- (d) the evaluation of ligand binding to an LBD without the need to use radiolabelled ligand derivatives or without reliance on the transcription factor properties of the corresponding nuclear receptor.
- (e) evaluating the effect of mutations in the LBD on ligand binding. Since SSR-LBD recombination requires ligand, mutations in the LBD that decrease ligand binding can be ascertained. Alternatively, mutations that improve binding of a different ligand can be selected. For example, ligand-dependent recombination could rearrange the DNA, so that an antibiotic resistance gene is expressed. Cells with the rearranged DNA will grow under the appropriate antibiotic selection, cells with unrearranged DNA will not. Specifically this describes a method for determining the effect of mutations in the LBD of a nuclear receptor on its ability to bind ligand, comprising the steps of (a) introducing mutations into the LBD of the SSR-LBD fusion protein, and (b) following steps (i) to (iii) above.

The invention also encompasses a method for regulating the recombination of DNA target sites, comprising the steps of:

According to the present invention, which contacting may be accomplished by direct introduction of the protein or by transformation of a cell with the nucleic acid encoding the protein.

(c) effecting a recombination of said DNA target sites by contacting the fusion protein with a ligand for the nuclear receptor component of the fusion protein, e.g. by adding the ligand to the system at a predetermined time, or by effecting production of the ligand in the system at a predetermined time.

#### Description of the Figures

Figure 1A depicts the plasmid, pNEO3GAL (thin line), integrated into the genome of 293 cells (thick line). FLP recombinase target sites (FRTs) are depicted as broad arrows and the neomycin resistance gene lies between the two FRTs. Upon estradiol administration, the DNA between the FRTs is excised, leaving one FRT in the genome and one on the excised circular DNA. The probe used for the Southern blot of Fig. 2 is also depicted.

Figure 1B depicts the steps involved in the creation of plasmid pHFE1.

Figure 2 shows a Southern blot of a timecourse of recombination occurring in the cell line P1.4. At the times indicated after estradiol, or ethanol only, administration in the upper part of the figure, cells were lysed, DNA purified, restricted with NdeI, run on a 1.25% agarose gel and blotted to Biodyne B membranes by standard methodologies. The membrane was hybridized with the radioactively labelled probe depicted in Fig. 1A. The upper band shows the unrecombined integrated DNA and the lower the recombined band. Estradiol was dissolved in ethanol and the extreme right hand lane shows cells treated with the equivalent amount of ethanol, without estradiol, for 51 hours.

Figure 3 shows the complete amino acid sequence of the SSR-LBD-

ding domain is located from amino acid 429-773.

recombinase domain, the linker peptide and the estrogen binding domain, respectively, are specified.

The following examples are offered by way of illustration and is not intended to limit the invention in any manner.

#### **Example 1**

##### **Construction of plasmid containing a SSR-LBD-fusion gene**

The experiment involves two plasmids, one of which is the target for recombination, and the other is the recombinase expression plasmid. As the target plasmid, pNEO $\beta$ GAL (9) was employed. pNEO $\beta$ GAL contains two FLP recombinase target sites (FRT) surrounding a gene encoding neomycin resistance (Fig. 1a). The expression plasmid, pHFE 1, carries two genes, one encoding an FLP recombinase-estrogen receptor LBD fusion protein and the other encoding hygromycin resistance (Fig. 1b). The plasmid was constructed by standard cloning procedures using pBluescribe (Stratagene), pOG 44 (9), pHE63 (22) and pCNH2 (27). The stop codon of the FLP recombinase coding region present in pOG44 was mutated by oligonucleotide replacement to introduce BamHI, BsiWI and EcoRI sites and continue the open reading frame. The BamHI site in the estrogen receptor encoded by pHE63 was used to join the coding regions of FLP recombinase and estrogen receptor LBD. All of the estrogen receptor coding region carboxy to the BamHI site is present in the plasmid, including its stop codon.

#### **Example 2**

##### **Regulation of site specific recombination**

pNEO $\beta$ GAL was introduced into 293 human embryonal kidney cells by electroporation ( $5 \times 10^6$  cells in 500  $\mu$ l phosphate-buffered

stant cells were cloned and characterised for incorporation of target plasmid DNA by Southern blotting. A clone, P1, showing

the second plasmid, pHFE 1 (1  $\mu$ g of XmnI linearised pHFE 1 was precipitated by the standard calcium phosphate method onto  $10^7$  P1 cells which were cultured in DMEM without phenol red, 10% charcoal stripped fetal calf serum). Four clones resistant to both neomycin and hygromycin (0.4 mg/ml G418, 0.4 mg/ml hygromycin B) were isolated and the dependance of recombination on estradiol administration was observed in all four. For clone P1.4, a time course of recombination in the presence of  $10^{-6}$  M estradiol is shown (Fig. 2).

### References

1. Falco, S.C., Li, Y., Broach, J.R. and Botstein, D. (1982) Cell 29, 573-584.
2. Cox, M.M. (1983) Proc. Natl. Acad. Sci. USA 80, 4223-4227.
3. Sauer, B. (1987) Mol. Cell. Biol. 7, 2087-2096.
4. Sauer, B. and Henderson, N. (1989) Nucleic Acids Res. 17, 147-161.
5. Golic, K.G. and Lindquist, S. (1989) Cell 59, 499-509.
6. Cregg, J.M. and Madden, K.R. (1989) Mol. Gen. Genet. 219, 320-323.
7. Matsuzaki, H. et al, (1990) J. Bacteriol. 172, 610-618.
8. Odell, J., Caimi, P., Sauer, B. and Russell, S. (1990) Mol. Gen. Genet. 223, 369-378.
9. O'Gorman, S., Fox, D.T. and Wahl, G.M. (1991) Science 251, 1351-1355.
10. Konsolaki, M. et al (1992) New Biologist 4, 551-557.
11. Lakso, M. et al (1992) Proc. Natl. Acad. Sci. USA 89, 6232-6236.
12. Baubonis, W. and Sauer, B. (1993) Nucleic Acids Res. 21, 2025-2029.
13. Stark, W.M. et al (1992) Trends Genet. 8, 432-439.
14. Utatsu, I. et al (1987) J. Bacteriol. 169, 5537-5545.
15. Chen, X.J. et al (1986) Nucleic Acids Res. 14, 4471-4481.
16. Klippel, A. et al (1993) EMBO J. 12, 1047-1057.

19. Webster, N.J.G. et al (1988) Cell 54, 194-207
20. Beato, M. (1989) Cell 56, 335-344.
21. Kumar, V. et al (1987) Cell 51, 941-951.
22. Picard, D., Salser, S.J. and Yamamoto, K.R. (1988) Cell 54, 1073-1080.
23. Eilers, M., Picard, D., Yamamoto, K.R. and Bishop, M. (1989) Nature 340, 66-68.
24. Superti-Furga, G. et al (1991) Proc. Natl. Acad. Sci. USA 88, 5114-5118.
25. Burk, O. and Klempenauer, K.-H. (1991) EMBO J. 10, 3713-3719.
26. Boehmelt, G. et al (1992) EMBO J. 11, 4641-4652.
27. Schmidt, E. et al (1990) Mol. Cell. Biol. 10, 4406-4411.



## Claims

1. A fusion protein, comprising a recombinase protein or a component of a recombinase complex, fused to part or all of a nuclear receptor so that the amino acids that bind the ligand of said nuclear receptor are included, such that, in cells or appropriate cell-free systems:
  - (a) recombinase activity is inhibited in the absence of ligand binding to said ligand binding domain, and
  - (b) recombinase activity is induced or altered by binding of ligand to said ligand binding domain.
2. The fusion protein of claim 1, wherein the nuclear receptor is a hormone receptor.
3. The fusion protein of claim 1 or 2, wherein the nuclear receptor is a steroid hormone receptor.
4. The fusion protein of any one of the claims 1-3, wherein the nuclear receptor is a mutated derivative of a nuclear receptor such that it retains the characteristics of the fusion protein of claim 1.
5. The fusion protein of any of the claims 1 - 4, wherein the nuclear receptor is a vertebrate glucocorticoid, estrogen, progesteron or androgen receptor.
6. The fusion protein of any one of the claims 1-5, wherein the recombinase protein or component of a recombinase complex is selected from the group consisting of: FLP recombinase, Cre recombinase, R recombinase from the Zygosaccharomyces rouxii plasmid pSR1, A recombinase from the Kluyveromyces drosophilarius plasmid pKD1, A recombinase
7. The fusion protein of any of claims 1-6,

wherein the recombinase protein or compound of a recombinase complex is fused to the nuclear receptor or ligand binding domain thereof by means of a genetic fusion.

8. The fusion protein of any of claims 1-7, wherein said nuclear receptor or ligand binding domain thereof is fused to the N- or C-terminus of said recombinase protein or component of a recombinase complex.
9. The fusion protein of any of claims 1-8, wherein said nuclear receptor is fused to said recombinase protein or component of a recombinase complex through a peptide linker.
10. The fusion protein of any of claims 1-9, which comprises the FLP recombinase and the ligand binding domain of the estrogen receptor.
11. A nucleic acid which encodes the fusion protein of any one of the claims 1-10.
12. The nucleic acid of claim 11, which is a DNA or RNA.
13. A recombinant vector comprising at least one copy of the nucleic acid of claims 11 or 12.
14. The vector of claim 13, which is a plasmid.
15. The plasmid pHFE1 (DSM 9265)
16. A cell containing a nucleic acid of claims 11 or 12 or a recombinant vector of any one of the claims 13-15.
17. A method for determining the binding of a ligand to the ligand binding domain of a nuclear receptor, comprising the

(a) the introduction of the fusion protein of claims 1-9, or the nucleic acid of claims 10 or 11 into cells, or appropriate cell-free systems, that contain the DNA target sites for recombination,

(b) optionally administering the ligand or a mixture suspected to contain a ligand or ligands, to be evaluated,

(c) detecting the recombinase activity, if any, of said fusion protein by detecting recombination or changes in the recombination rate between said DNA target sites.

19. The method of claim 18,  
wherein the recombination between said DNA target sites is detected by direct measurement of the physical arrangement of said target sites.
20. The method of claim 18,  
wherein the recombination between said DNA target sites is detected by assessing the properties encoded by the DNA regions carrying the target sites before or after recombination.
21. A method for determining the effect of mutations in the ligand binding domain of a nuclear receptor on its ability to bind ligand, comprising the steps of:
  - (a) introducing mutations into the ligand binding domain of the fusion protein of claims 1-10,
  - (b) following steps (a) to (c) of claim 18.
22. A method for regulating the recombination of DNA target sites, comprising the steps of:
  - (a) providing cells or appropriate cell-free systems that contain DNA target sites for a site specific recombinase,

tacting the fusion protein with a ligand for the nuclear receptor component of said fusion protein.

FIG. 1A




FIG. 1B




FIG. 2




Fig. 3

FLP/EBD AMINO ACID SEQUENCE AS ENCODED BY PHFE1

1 MPQFDILCKT PPKVLVRQFV ERFERPSGEK IALCAAELTY LCWMITHNGT  
51 AIKRATFMSY NTIISNSLSL DIVNKSLOFK YKTQKATILE ASLKKLIPAW  
101 EFTIIPYYGQ KHQSDITDIV SSLQLQFESS EEADKGNHS KSMLKALLSE  
151 GESTWEITEK ILNSFEYTSR FTKTKTLYQF LFLATFINCG RFSDIKNVDP  
201 KSFKLVQNKY LGVIIQCLVT ETKTSVSRHI YFFSARGRID PLVYLDEFLR  
251 NSEPVLKRVN RTGNSSSNKQ EYQLLDNLV RSYNKALKKN APYSIFAIKN  
301 GPKSHIGRHL MTSFLSMKGL TELTNVVGWV SDKRASAVAR TTYTHQITAI  
351 PDHYFALVSR YYAYDPISKE MIALKDETNP IEEWQHIEQL KGSAEGSIRY  
401 PAWNGHISQE VLDYLSSYN RRI

- FLP ENDS HERE -

424 SVRGS

- LINKER PEPTIDE ENDS HERE -

429 MK GGIRKDRRG RMLKHKRQRD

451 DGEGRGEVGS AGDMRAANLW PSPLMIKRSK KNSLALSITA DQMVSALLDA

501 EPPILYSEYD PTRPFSEASM MGLLTNLADR ELVHMINWAK RVPGFVDLTL

551 HDQVHILLECA WLEILMIGLV WRSMEHPVKL LFAPNLLDR NQGKCVEGMV

601 EIFDMLLATS SRFRMMNLQG EEFVCLKSII LLNSGVYTFL SSTLKSLEEK

651 DHIHRVLDKI TDTLIHLMAG AGLTLQQQHQLAQLLLILS HIRHMSNKG

- ESTROGEN BINDING DOMAIN ENDS HERE -

Fig. 4/1

CODING SEQUENCE OF THE FLP/EBD FUSION PROTEIN ENCODED BY  
 PHFE1

ATGCCACAATTTGATATATTATGTAAAACACCACCTAAGGTGCTTGTTTCGT  
CAGTTTGTGGAAAGGTTTGAAAGACCTTCAGGTGAGAAAATAGCATTATG  
TGCTGCTGAACCTAATTTATGTTGGATGATTACACATAACGGAACAGC  
AATCAAGAGAGCCACATTCATGAGCTATAATACTATCATAAGCAATTTCG  
TGAGTTCCGATATTGTCAACAAGTCACTGCAGTTTAAATACAAGACGCAA  
AAAGCAACAATTCTGGAAGCCTCATTAAAGAAATTGATTCCTGCTTGGA  
ATTTACAATTATTCCITACTATGGACAAAAACATCAATCTGATATCACTG  
ATATTGTAAGTAGTTTGCAATTACAGTTCGAATCATCGGAAGAAGCAGAT  
AAGGGAAATAGCCACAGTAAAAAAATGCTTAAAGCACTTCTAAGTGAGG  
GTGAAAGCATCTGGGAGATCACTGAGAAAATACTAAATTCGTTTGAGTAT  
ACTTCGAGATTTACAAAAACAAAACTTTATACCAATTCCTCTTCCTAGC  
TACTTTTCATCAATTGTGGAAGATTCAGCGATATTAAGAACGTTGATCCGA  
AATCATTTAAATTAGTCCAAAATAAGTATCTGGGAGTAATAATCCAGTG  
TTTAGTGACAGAGACAAAGACAAGCGTTAGTAGGCACATATACTTCTTTA  
GCGCAAGGGGTAGGATCGATCCACTTGTATATTTGGATGAATTTTTGAGGA  
ATTCTGAACCAAGTCCTAAAACGAGTAAATAGGACCGCAATTCTTCAAGC  
AACAAGCAGGAATACCAATTATTA AAAAGATAACTTAGTCAGATCGTACA  
ACAAAGCTTTGAAGAAAAATGCGCCTTATTCAATCTTTGCTATAAAAAA  
TGGCCCAAAATCTCACATTGGAAGACATTTGATGACCTCATTCTTTCAAT  
GAAGGGCCTAACGGAGTTGACTAATGTTGTGGGAAATTGGAGCGATAAGCG  
TGCTTCTGCCGTGGCCAGGACAACGTATACTCATCAGATAACAGCAATACCT  
GATCACTACTTCGCACTAGTTTCTCGGTACTATGCATATGATCCAATATCA  
AAGGAAATGATAGCATTGAAGGATGAGACTAATCCAATTGAGGAGTGGC  
AGCATATAGAACAGCTAAAGGGTAGTGCTGAAGGAAGCATACGATACCCC  
GCATGGAATGGGATAATATCACAGGAGGTACTAGACTACCTTTCATCCTAC  
ATAAATAGACGCATA

-FLP ENDS HERE-

TCCGTACGCGGATCC

- SYNTHETIC LINKER SEQUENCE ENDS HERE -

ATGAAAGGTGGGATACGAAAAGACCGAAGAGGAGGGAGAATGTTGAAAC  
ACAAGCGCCAGAGAGATGATGGGGAGGGCAGGGGTGAAGTGGGGTCTGCTG

ATCAGTGAAGGCTTCGATGATGGGCTTACGACCAACCTTAACTCTTCTTAACT  
TGGTTCACATGATCAACTGGGCGAAGAGGGTGCCAGGCTTTGTGGATTTGAC



Fig. 4/2

CCTCCATGATCAGGTCCACCTTCTAGAATGTGCCTGGCTAGAGATCCTGATG  
ATTGGTCTCGTCTGGCGCTCCATGGAGCACCCAGTGAAGCTACTGTTTGCTCCT  
AACTTGCTCTTGGACAGGAACCAGGGAAAATGTGTAGAGGGCATGGTGGAG  
ATCTTCGACATGCTGCTGGCTACATCATCTCGGTTCCGCATGATGAATCTGCA  
GGGAGAGGAGTTTGTGTGCCTCAAATCTATTATTTTGCTTAATTCTGGAGTG  
TACACATTTCTGTCCAGCACCCCTGAAGTCTCTGGAAGAGAAGGACCATATCC  
ACCGAGTCCTGGACAAGATCACAGACACTTTGATCCACCTGATGGCCAAGGC  
AGGCCTGACCCTGCAGCAGCAGCACCAGCGGCTGGCCCAGCTCCTCCTCATCCT  
CTCCCACATCAGGCACATGAGTAACAAAGGCATGGAGCATCTGTACAGCAT  
GAAGTGCAAGAACGTGGTGCCCCTCTATGACCTGCTGCTGGAGATGCTGGAC  
GCCCACCGCCTACATGCGCCCACTAGCCGTGGAGGGGCATCCGTGGAGGAGAC  
GGACCAAAGCCACTTGGCCACTGCGGGCTCTACTTCATCGCATTCTTGCAAA  
AGTATTACATCACGGGGGAGGCAGAGGGTTCCCTGCCACAGTCTGA

- HORMONE BINDING DOMAIN ENDS HERE -

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 94/02088

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 5 C07K15/04 C12N15/62

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	THE EMBO JOURNAL, vol.11, no.12, 1992, OXFORD pages 4641 - 4652 BOEHMELT ET AL. 'Hormone-regulated v-rel estrogen receptor fusion protein: ...' cited in the application see the whole document ---	1-6, 10-12, 16,17
A	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, vol.88, no.12, 15 June 1991, WASHINGTON, DC pages 5114 - 5118 SUPERTI-FURGA ET AL. 'Hormone-dependent transcriptional regulation .....' cited in the application see abstract --- -/--	1-6, 10-12, 16,17

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

14 October 1994

Name and mailing address of the ISA  
European Patent Office, P.B. 5818 Patentaan 2  
NL - 2280 HV Rijswijk  
Tel. (31-70) 340-2040, 1-  
(31-70) 340-3014

Authorized officer

EDU

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 94/02088

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CELL, vol.54, 23 September 1988, CAMBRIDGE, MASS. pages 1073 - 1080 PICARD ET AL. 'A movable and regulable inactivation function ..... <p>-----</p> <p>cited in the application see abstract</p>	1
A	EP,A,0 146 050 (CALLEWAERT) 26 June 1985 see abstract	1
A	NUCLEIC ACIDS RESEARCH, vol.14, no.11, 1986, LONDON pages 4471 - 4481 CHEN ET AL. 'Sequence organization of the circular plasmid pKD1 ..... <p>-----</p> <p>cited in the application see abstract</p>	2-6,10
A	NUCLEIC ACIDS RESEARCH, vol.17, no.1, 1989, LONDON pages 147 - 161 SAUER ET AL. 'Cre-stimulated recombination .....' <p>-----</p> <p>cited in the application see abstract</p>	2-6,10
A	TRENDS IN GENETICS, vol.8, 1992, AMSTERDAM pages 432 - 439 STARK ET AL. 'Catalysis by site-specific recombinases' <p>-----</p> <p>cited in the application</p>	

### Information on patent family members

PCT/EP 94/02088

Form PCT ISA 2.0 (patent family annex) (July 1992)